

REPRESENTATION-FINITE TRIVIAL EXTENSION ALGEBRAS

Ibrahim ASSEM

*Department of Mathematics, University of Ottawa, 585, King Edward, Ottawa, Ont., K1N 9B4
Canada*

Dieter HAPPEL

Fakulat für Mathematik, Universität Bielefeld, D-4800 Bielefeld, Fed. Rep. Germany

Oscar ROLDÁN

*Escuela de Matematica, Universidad de Costa Rica, Ciudad Universitaria Rodrigo Facio, San
José, Costa Rica*

Communicated by H. Bass

Received 12 October 1983

Let A be a finite-dimensional basic connected associative algebra over an algebraically closed field, and $T(A) = A \ltimes DA$ its trivial extension by its minimal injective cogenerator. We prove that $T(A)$ is representation-finite of Cartan class Δ if and only if A is an iterated tilted algebra of Dynkin class Δ . The proof also yields a construction procedure for iterated tilted algebras of Dynkin type.

Introduction

Let A be a finite-dimensional basic connected algebra over an algebraically closed field, and denote by $T(A) = A \ltimes DA$ the trivial extension of A by its minimal injective cogenerator DA . Then $T(A)$ is self-injective and, in fact, symmetric. It is thus natural to ask for a description of the algebras A such that $T(A)$ is representation-finite. This problem was already considered by Müller [15], Green and Reiten [9], and Iwanaga and Wakamatsu [14] in the case where the square radical of A is equal to zero. It was shown by Tachikawa [18] that if A is hereditary and representation-finite, then $T(A)$ is again representation-finite (see also [19]). In [20], Yamagata proved that if $T(A)$ is representation-finite, then the ordinary quiver of A contains no oriented cycles. Later, Hughes and Waschbüsch [13] and Hoshino [12] proved that if A is a tilted algebra of Dynkin type Δ , then $T(A)$ is representation-finite of Cartan class Δ , and conversely, if $T(A)$ is representation-finite of Cartan class Δ , then there exists a tilted algebra B of Dynkin type Δ such that $T(A) \cong B \ltimes DB$ (compare also [7], [17]). The purpose of the present paper is to prove the following theorem:

Theorem. *Let A be a finite-dimensional basic connected k -algebra and $T(A) = A \ltimes DA$ be its trivial extension. Then $T(A)$ is representation-finite of Cartan class Δ if and only if A is an iterated tilted algebra of Dynkin type Δ .*

This result was already obtained in the case where $\Delta = \mathbb{A}_n$ in [13], using the classification of the iterated tilted algebras of type \mathbb{A}_n given in [1]. The proof presented here, however, does not use the classification results for iterated tilted algebras of Dynkin type. As a corollary, we obtain a simple construction procedure for all iterated tilted algebras of Dynkin type Δ starting from the tilted algebras of the same type.

1. Preliminaries

1.1. We shall let k denote a fixed algebraically closed field, and A a basic, connected, finite-dimensional k -algebra. By $\text{mod } A$ will be meant the category of finite-dimensional right A -modules. The letter Q will be used to denote the ordinary quiver of A , and i, j, \dots to denote its vertices. The simple A -module corresponding to the vertex i of Q will be denoted by $S(i)$, and its projective cover (respectively, its injective hull) by $P(i)$ (respectively, $I(i)$). We shall freely use properties of the Auslander-Reiten sequences and the Auslander-Reiten quiver, for both finite-dimensional and locally finite-dimensional algebras, and refer for these to [8].

1.2. A module T_A is called a tilting module [10] if $\text{Ext}_A^1(T, T) = 0$, $\text{Ext}_A^2(T, -) = 0$ and there exists a short exact sequence $0 \rightarrow A_A \rightarrow T'_A \rightarrow T''_A \rightarrow 0$, with T' and T'' direct sums of summands of T .

A finite-dimensional k -algebra B is called iterated tilted of type Δ [1] if there exists a sequence of algebras $A_0, A_1, \dots, A_m = B$ where A_0 is a hereditary algebra having Δ as underlying graph of its ordinary quiver, and a sequence of tilting modules $(T_{A_i}^{(i)})_{0 \leq i \leq m-1}$ such that $\text{End } T_{A_i}^{(i)} = A_{i+1}$ and, for every indecomposable A_{i+1} -module M , we have either $M \otimes_{A_{i+1}} T^{(i)} = 0$, or $\text{Tor}_1^{A_{i+1}}(M, T^{(i)}) = 0$. If $m \leq 1$, B is called a tilted algebra [10]. The iterated tilted algebras of type \mathbb{A}_n were classified in [1]. A characterisation of the iterated tilted algebras of Dynkin type is given in [11].

1.3. Let $D = \text{Hom}_k(-, k)$ denote the standard duality of $\text{mod } A$. The trivial extension of A by its minimal injective cogenerator ${}_A(DA)_A$ is defined to be the algebra $T = A \ltimes DA$ whose additive structure is that of the group $A \oplus DA$, with the multiplication defined by

$$(x, f)(y, g) = (xy, xg + fy)$$

for $x, y \in A$ and $f, g \in {}_A(DA)_A$. It is well-known that the trivial extension algebra T is self-injective, and in fact symmetric. Thus, a connected trivial extension algebra

T is representation-finite if and only if the stable part of its Auslander–Reiten quiver is isomorphic to $\mathbb{Z}\Delta/G$, where Δ is a Dynkin diagram (the Cartan class of T), and G an admissible group of automorphisms of $\mathbb{Z}\Delta$ [16].

1.4. Consider the matrix algebra:

$$\hat{A} = \begin{bmatrix} \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \\ & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \\ & & M_{n-1} & A_{m-1} & & & & \\ & & & M_m & A_m & & & \\ & & & & M_{m+1} & A_{m+1} & & \\ & & & & & \ddots & \ddots & \ddots \end{bmatrix}$$

in which matrices have only finitely many non-zero entries, $A_m = A$, $M_m = {}_A(DA)_A$ for all $m \in \mathbb{Z}$, all the remaining entries are zero, and multiplication is induced from the canonical maps $A \otimes_A DA \rightarrow DA$, $DA \otimes_A A \rightarrow DA$ and the zero map $DA \otimes_A DA \rightarrow 0$ [13]. Clearly, \hat{A} is locally finite-dimensional [6]. The identity maps $A_m \rightarrow A_{m-1}$, $M_m \rightarrow M_{m-1}$ induce an automorphism ν of \hat{A} , and the orbit space \hat{A}/ν inherits the structure of a finite-dimensional k -algebra, isomorphic to $T = A \ltimes DA$. Moreover, ν induces an automorphism of the category $\text{mod } \hat{A}$ of finite-dimensional \hat{A} -modules, and the quotient of the Auslander–Reiten quiver $\Gamma_{\hat{A}}$ under the automorphism ν is a set of complete connected components of Γ_T [13].

Observe that any complete set $(e_i)_{1 \leq i \leq n}$ of primitive orthogonal idempotents of A can be lifted to a complete set of primitive orthogonal idempotents $(e_{(i,m)})_{1 \leq i \leq n, m \in \mathbb{Z}}$ of \hat{A} . Thus ν induces also an automorphism of the ordinary quiver \hat{Q} of \hat{A} . A full connected subquiver Q' of \hat{Q} is called a complete ν -slice of \hat{Q} if its vertex set Q'_0 consists of a complete set of representatives of the ν -orbits of \hat{Q}_0 , and for any path $i = i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_t = j$ such that $i, j \in Q'_0$, we have $i_s \in Q'_0$ for all $0 < s < t$. Given a complete ν -slice Q' of \hat{Q} , the algebra

$$\text{End}_{\hat{A}} \left(\bigoplus_{(i,m) \in Q'_0} e_{(i,m)} \hat{A} \right)$$

is called the algebra of Q' , and denoted by $\text{Alg}(Q')$. Then, if the quiver Q of A has no oriented cycles, and B is a finite-dimensional k -algebra, the following assertions are equivalent [13]:

- (i) $B \ltimes DB \xrightarrow{\sim} A \ltimes DA$,
- (ii) $\hat{B} \xrightarrow{\sim} \hat{A}$,
- (iii) $B \xrightarrow{\sim} \text{Alg}(Q')$ for some complete ν -slice Q' of \hat{Q} .

1.5. For a sink i in the quiver of A , we define the ν -reflection of Q , $\sigma_i^+ Q$ to be the full subquiver of \hat{Q} with vertex set $\{(j, 0) \in \hat{Q}_0 \mid j \in Q_0, j \neq i\} \cup \{(i, 1)\}$. A ν -reflection sequence of sinks i_1, i_2, \dots, i_t is a sequence of vertices of \hat{Q} such that i_s is a sink of $\sigma_{i_{s-1}}^+ \dots \sigma_{i_1}^+ Q$ for $1 \leq s \leq t$. ν -reflection sequences of sources are defined dually. Then the complete ν -slices of \hat{Q} are exactly the repeated ν -reflections of Q [13].

For a sink i of Q , let $T_i^+ A$ denote the one-point extension algebra

$$T_i^+A = \begin{bmatrix} A & 0 \\ I(i)_A & k_i \end{bmatrix}$$

(where $k_i = \text{End } S(i) \curvearrowright k$), and S_i^+A be the quotient of T_i^+A by the two-sided ideal generated by the idempotent e_i corresponding to i . Then T_i^+A has a complete set $e_1, e_2, \dots, e_n, e_i$ of $n + 1$ primitive orthogonal idempotents, and the ordinary quiver of S_i^+A is obtained from that of T_i^+A by deleting the vertex corresponding to e_i . Similarly, starting with a source j of Q , we may define the algebras T_j^-A and S_j^-A . Also, $S_i^+A \curvearrowright \text{Alg}(\sigma_i^+Q)$ and $S_j^-A \curvearrowright \text{Alg}(\sigma_j^-Q)$ [13], thus we call S_i^+A and S_j^-A the reflections of A .

2. Reflections of iterated tilted algebras

2.1. We shall first recall some definitions. Let A be an algebra of finite global dimension, the Cartan matrix of A is defined to be the $n \times n$ matrix $C_A = [c_{ij}]$, where

$$c_{ij} = \dim_k \text{Hom}_A(P(i), P(j))$$

and the Euler characteristic of A is defined to be the bilinear form on the Grothendieck group $K_0(A)$ of A with matrix $B_A = [b_{ij}]$ given by

$$b_{ij} = \sum_{l \geq 0} (-1)^l \dim_k \text{Ext}_A^l(S(i), S(j))$$

(this sum is in fact finite due to our hypothesis on A). We shall let χ_A denote the associated quadratic form. It is easily seen that C_A is invertible and that its inverse equals the transpose B_A^t of B_A . We have the following lemma:

Lemma. *Let A be an algebra such that the quiver Q of A has no oriented cycles, and i be a sink of Q . Then the Cartan matrices of A and $A' = S_i^+A$ are congruent, that is, there exists a matrix $X \in \text{GL}(n, \mathbb{Z})$ such that $C_{A'} = XC_A X^t$.*

Proof. We may assume, without loss of generality, that $i = 1$. Observe that, since Q has no oriented cycles, the global dimension of A is finite. Let $1'$ denote the vertex added to Q in the construction of T_1^+A and let $P'(1'), P'(2), \dots, P'(n)$ denote the non-isomorphic indecomposable projective A' -modules. By construction, we have $\text{rad } P'(1')_{A'} = I(1)_A / \text{soc } I(1)$.

If $i \neq 1, j \neq 1$, we have

$$\dim_k \text{Hom}_{A'}(P'(i), P'(j)) = \dim_k \text{Hom}_A(P(i), P(j)).$$

It is clear that

$$\dim_k \text{Hom}_{A'}(P'(1'), P'(1')) = 1 = \dim_k \text{Hom}_A(P(1), P(1)).$$

If $i \neq 1, i' \neq 1'$, we have

$$\mathrm{Hom}_{A'}(P'(1'), P'(i)) = 0 = \mathrm{Hom}_A(P(i), P(1)).$$

Finally, if $i \neq 1$, $i \neq 1'$, we have

$$\begin{aligned} \dim_k \mathrm{Hom}_{A'}(P'(i), P'(1')) &= \dim_k \mathrm{Hom}_A(P(i), I(1)) \\ &= \dim_k \mathrm{Hom}_A(P(1), P(i)). \end{aligned}$$

Hence, let $u = -\dim I(1)_A$ and consider the matrix $X = [u \ e_2 \cdots e_n]$ with u as first column, and e_i ($2 \leq i \leq n$) the canonical basis vectors. Then $X \in \mathrm{GL}(n, \mathbb{Z})$ (in fact, X^2 is the identity matrix), and a straightforward calculation shows that $C_{A'} = XC_A X^t$.

2.2. Proposition. *Let A be an iterated tilted algebra of Dynkin type Δ , and i be a sink in the quiver Q of A . Then $S_i^+ A$ is also iterated tilted of type Δ .*

Proof. By [13], $S_i^+ A$ has a preprojective component \mathcal{C} in its Auslander-Reiten quiver. If M is an indecomposable module in \mathcal{C} , then its dimension-vector $\dim M$ is a root of the quadratic form $\chi_{S_i^+ A}$. Since the previous lemma shows that $\chi_{S_i^+ A}$ is congruent to χ_A , and by [10], χ_A is congruent to the Tits form of a hereditary algebra of Dynkin type Δ , it follows that $\chi_{S_i^+ A}$ has only finitely many roots. Hence \mathcal{C} is a finite component of $\Gamma_{S_i^+ A}$ and, by a theorem of Auslander [4], $S_i^+ A$ is representation-finite. In fact, $S_i^+ A$ is simply connected, since it satisfies the (S)-condition [5] (for, A being simply connected [2], the indecomposable injective A -module $I(i)_A$ has separated socle factor, hence the indecomposable projective $S_i^+ A$ -module $P'(i')$ has separated radical. It is clear that the other indecomposable projective $S_i^+ A$ -modules have separated radicals). Now, it is proved in [11] that a simply connected algebra is iterated tilted of Dynkin type Δ provided the quadratic form associated to its Euler characteristic is congruent to the Tits form of a hereditary algebra of Dynkin type Δ . This shows that $S_i^+ A$ is indeed iterated tilted of Dynkin type Δ .

Dually, if j is a source of Q , $S_j^- A$ is also iterated tilted of type Δ . Inductively, if i_1, i_2, \dots, i_t is a v -reflection sequence of sinks (respectively, if j_1, j_2, \dots, j_s is a v -reflection sequence of sources), then $S_{i_t}^+ \cdots S_{i_1}^+ A$ (respectively, $S_{j_s}^- \cdots S_{j_1}^- A$) is also iterated tilted of type Δ .

3. The main result

3.1. Theorem. *Let A be a basic, connected finite-dimensional k -algebra, and let $T = A \ltimes DA$ be its trivial extension by its minimal injective cogenerator. Then T is representation-finite of Cartan class Δ if and only if A is an iterated tilted algebra of Dynkin type Δ .*

Proof. Let A be an iterated tilted algebra of Dynkin type Δ , we shall prove that

$T = A \ltimes DA$ is representation-finite of type Δ . Since A is simply connected [2], its ordinary quiver Q must contain at least one strong sink i (that is to say, a vertex $i \in Q_0$ such that there is no chain of indecomposable modules and irreducible maps $I(j)_A \rightarrow \cdots \rightarrow I(i)_A$ for any $j \in Q_0$). A strong sink is always a sink [13].

Clearly, there are canonical embeddings of $\text{mod } A$ and $\text{mod}(S_i^+ A)$ into $\text{mod}(T_i^+ A)$, where the first category can be described as the class of $T_i^+ A$ -modules without the simple injective $I(i')_{T_i^+ A}$ as a composition factor, while the second is the class of modules without the simple projective $P(i)_{T_i^+ A}$ as a composition factor. Moreover, the only $T_i^+ A$ -module which is not in either one of these classes is the projective-injective module $P(i')_{T_i^+ A} = I(i)_{T_i^+ A}$ [13]. Since, by Proposition 2.2, $S_i^+ A$ is also iterated tilted of type Δ , we may iterate this process to obtain the Auslander-Reiten quiver of \hat{A} . Let i_1, i_2, \dots, i_t be a strong ν -reflection sequence of sinks, that is to say, a sequence such that i_r is a strong sink in the quiver $\sigma_{i_{r-1}}^+ \cdots \sigma_{i_1}^+ Q$ of the algebra $S_{i_{r-1}}^+ \cdots S_{i_1}^+ A$ (for all $1 \leq r \leq t$), then the algebra $S_{i_t}^+ \cdots S_{i_1}^+ A$ is iterated tilted of type Δ , and a straightforward induction yields the Auslander-Reiten quiver of $T_{i_t}^+ \cdots T_{i_1}^+ A$. Again, any indecomposable $T_{i_t}^+ \cdots T_{i_1}^+ A$ -module is either projective-injective, or else can be identified with an indecomposable module over one of the iterated tilted algebras $S_{i_r}^+ \cdots S_{i_1}^+ A$ ($1 \leq r \leq t$). This procedure yields in fact a connected component Γ of the Auslander-Reiten quiver of \hat{A} . Indeed, we construct inductively strong ν -reflection sequence of sinks $(i_t)_{t \geq 1}$ and of sources $(j_s)_{s \geq 1}$, then every vertex of \hat{Q} belongs to $\bigcup_{t=0}^{\infty} \sigma_{i_t}^+ \cdots \sigma_{i_1}^+ Q$ or to $\bigcup_{s=0}^{\infty} \sigma_{j_s}^- \cdots \sigma_{j_1}^- Q$ and the categories $\text{mod}(T_{i_t}^+ \cdots T_{i_1}^+ A)$, and $\text{mod}(T_{j_s}^- \cdots T_{j_1}^- A)$ embed naturally in $\text{mod } \hat{A}$. Thus, as in [13], we obtain a connected component Γ of $\Gamma_{\hat{A}}$, containing all projective \hat{A} -modules and in which all modules have bounded composition length. It then follows from [4] that $\Gamma = \Gamma_{\hat{A}}$. On the other hand, the orbit space $\Gamma_{\hat{A}}/\nu$ is a connected component of Γ_T , which contains all the projective T -modules, and in which all modules have bounded length. Therefore $\Gamma_{\hat{A}}/\nu$ is actually equal to Γ_T , and T is representation-finite.

We next show that the Cartan class Δ' of T is equal to Δ . Indeed, T being a representation-finite trivial extension algebra, there exists a tilted algebra B of type Δ' such that $T \cong B \ltimes DB$ ([12], [13]). However, $A \ltimes DA \cong B \ltimes DB$ implies that B is the algebra of a complete ν -slice Q' of \hat{Q} , which must be in fact a repeated ν -reflection of Q . More precisely, there exists a ν -reflection sequence of sinks i_1, i_2, \dots, i_t (or of sources j_1, j_2, \dots, j_s) of Q such that $Q' = \sigma_{i_t}^+ \cdots \sigma_{i_1}^+ Q$ resp., $Q' = \sigma_{j_s}^- \cdots \sigma_{j_1}^- Q$ and therefore $B \cong S_{i_t}^+ \cdots S_{i_1}^+ A$ (resp., $B \cong S_{j_s}^- \cdots S_{j_1}^- A$). This indeed follows by induction on t (resp., s) and the fact that, for a sink i , (resp., a source j) of Q , $S_i^+ A \cong \text{Alg}(\sigma_i^+ Q)$ (resp., $S_j^- A \cong \text{Alg}(\sigma_j^- Q)$). However, A being iterated tilted of type Δ , Proposition 2.2 implies that B is of the same type, hence $\Delta = \Delta'$.

We have thus proved that if A is iterated tilted of Dynkin type Δ , its trivial extension algebra $A \ltimes DA$ is representation-finite of Cartan class Δ . Conversely, if $A \ltimes DA$ is representation-finite of Cartan class Δ , there exists a tilted algebra B of type Δ such that $B \ltimes DB \cong A \ltimes DA$ [13]. Using the same argument as above, we can find a ν -reflection sequence of sinks i_1, i_2, \dots, i_t (or of sources j_1, j_2, \dots, j_s) of

Q such that $A \xrightarrow{\sim} S_{i_l}^+ \cdots S_{i_1}^+ B$ (resp., $A \xrightarrow{\sim} S_{j_1}^- \cdots S_{j_k}^- B$). Applying again Proposition 2.2, it follows that A is iterated tilted of type Δ .

3.2. We have the following obvious corollary:

Corollary. *Let A be an iterated tilted algebra of Dynkin type Δ . Then there exists a tilted algebra B of type Δ , and a v -reflection sequence of sinks i_1, i_2, \dots, i_l in the quiver of B such that $A \xrightarrow{\sim} S_{i_l}^+ \cdots S_{i_1}^+ B$.*

3.3. Remark. Using the classification of the iterated tilted algebras of types \mathbb{B}_n and \mathbb{C}_n [2], it is shown in [3] that the main theorem holds in these cases too. Finally, an easy computation shows that it holds as well in the cases \mathbb{F}_4 and \mathbb{G}_2 .

Acknowledgement

This research was completed while the second author was visiting the Universities of Carleton and Ottawa, for which he would like to thank Professor V. Dlab for his support.

References

- [1] I. Assem and D. Happel, Generalized tilted algebras of type \mathbb{A}_n , *Comm. Algebra* 9 (1981) 2101–2125.
- [2] I. Assem, Iterated tilted algebras of types \mathbb{B}_n and \mathbb{C}_n , *J. Algebra* 84 (2) (1983) 361–390.
- [3] I. Assem and O. Roldán, Gradings of \mathbb{B}_n and \mathbb{C}_n of finite representation type, *Trans. Amer. Math. Soc.* 279 (2) (1983) 589–609.
- [4] M. Auslander, Applications of morphisms determined by objects, *Proc. Conf. on Representation Theory (Philadelphia 1976)* (Marcel Dekker, New York, 1978), 245–327.
- [5] R. Bautista, F. Larrión and L. Salmerón, On simply connected algebras, *J. London Math. Soc.* 27 (2) (1983) 212–220.
- [6] K. Bongartz and P. Gabriel, Covering spaces in Representation Theory, *Invent. Math.* 65 (1982) 331–378.
- [7] O. Bretscher, C. Läser and C. Riedtmann, Selfinjective and simply connected algebras, *Manuscripta Math.* 36 (3) (1981) 253–308.
- [8] P. Gabriel, Auslander–Reiten sequences and representation-finite algebras, *Proc. ICRA II (1979)*, *Lecture Notes in Math.* 831 (Springer, Berlin, 1980) 1–71.
- [9] E.L. Green and I. Reiten, On the construction of ring extensions, *Glasgow Mat. J.* 17 (1976) 1–11.
- [10] D. Happel and C.M. Ringel, Tilted algebras, *Trans. Amer. Math. Soc.* 274 (2) (1982) 399–443.
- [11] D. Happel, Tilting sets on cylinders, in preparation.
- [12] M. Hoshino, Trivial extensions of tilted algebras, *Comm. Algebra* 10 (18) (1982) 1965–1999.
- [13] D. Hughes and J. Waschbüsch, Trivial extensions of tilted algebras, *Proc. London Math. Soc.* 46 (3) (1983) 347–364.
- [14] Y. Iwanaga and T. Wakamatsu, Trivial extensions of artin algebras, *Proc. ICRA II (1979)*, *Lecture Notes in Math.* 832 (Springer, Berlin, 1980) 295–301.
- [15] W. Müller, Unzerlegbare Moduln über artinschen Ringen, *Math. Z.* 137 (1974) 197–226.

- [16] C. Riedtmann, Algebren, Darstellungsköcher, Ueberlagerungen und Zurück, *Comment. Math. Helv.* 55 (1980) 199–224.
- [17] C.M. Ringel, Tilted algebras and self-injective algebras, Preprint, University of Bielefeld (1981).
- [18] H. Tachikawa, Representations of trivial extensions of hereditary algebras, *Proc. ICRA II* (1979), *Lecture Notes in Math.* 832 (Springer, Berlin, 1980) 579–599.
- [19] K. Yamagata, Extensions over hereditary artinian rings with self-dualities I, *J. Algebra* 73 (2) (1981) 386–433.
- [20] K. Yamagata, On algebras whose trivial extensions are of finite representation type, *Proc. ICRA III* (1980), *Lecture Notes in Math.* 903 (Springer, Berlin, 1981) 364–371.